


Central Venous Catheter Dressing Change Algorithm for Adults



- Part #s**
- Tincture green soap (PS#991453)
 - Triple povidone iodine swabs (PS#907171)
 - Transparent tegaderm (PS# 51453)
 - Sorbaview (PS#924250)
 - IV 3000 tegaderm (PS# 116065)
 - Adult central line drsg tray w/CHG w/ device (PS#98287)
 - CHG drsg change kit tray, ports (PS#1062)
 - Drsg change kit tray, ports w/out CHG (PS# 65423)
 - Drsg PICC/CVC w/CHG and device (PS#96740)
 - Biopatch (PS#92687)
 - StatSeal powder (PS#122133)
 - StatSeal disc (PS# 122134)
 - CHG tegaderm drsg (PS#21049)
 - Shoulder length glove (PS#103103)
 - Adhesive remover wipes (PS# 98966)
 - Adhesive remover spray (PS# 98965)
 - CHG tegaderm, large lumen (PS#21049)
 - Drape (PS# 907043)

- Other Recommendations:**
- Adhesive remover spray/wipes may be used to loosen the dressing for patients with poor skin integrity or at risk of poor skin integrity
 - Allow skin to dry completely to prevent skin damage, may take 3-5min.
 - Consult wound/ostomy nursing
 - If patients require frequent dressing change (>3/24h) consider changing to another type dressing Barrier film is recommended for all changes, found in dressing change kit
 - StatSeal for hemostasis by trained RN's BMT/GVH/Radiation/Thiotepa patients- may use dry dressing (betadine, securement device, sterile gauze, burn net, changed q24h and PRN)

* 7 Days or as needed (damp, loosened, visbly soiled, or skin change issues)